

# 2i

## Infection et Inflammation



## IDENTIFICATION OF VIRULENCE MARKERS OF MYCOBACTERIUM ABSCESSUS FOR INTRACELLULAR REPLICATION IN PHAGOCYTES

**What differentiates *Mycobacterium abscessus* from other saprophytic mycobacteria is the ability to resist phagocytosis by human macrophages and the ability to multiply inside such cells. Fabienne Mishguich explains us how and why.**

These virulence traits render *M. abscessus* pathogenic, especially in vulnerable hosts with underlying structural lung disease, such as cystic fibrosis, bronchiectasis or tuberculosis. How patients become infected with *M. abscessus* remains unclear. Unlike many mycobacteria, *M. abscessus* is not found in the environment but might reside inside amoebae, environmental phagocytes that represent a potential reservoir for *M. abscessus*. Indeed, *M. abscessus* is resistant to amoebal phagocytosis and the intra-amoeba life seems to increase *M. abscessus* virulence in an experimental model of infection. However, little is known about *M. abscessus* virulence in itself. To decipher the

genes conferring an advantage to *M. abscessus* intracellular life, a screening of a *M. abscessus* transposon mutant library was developed. In parallel, a method of RNA extraction from intracellular *Mycobacteria* after co-culture with amoebae was developed. This method was validated and allowed the sequencing of whole *M. abscessus* transcriptomes inside the cells; providing, for the first time, a global view on *M. abscessus* adaptation to intracellular life. Both approaches give us an insight into *M. abscessus* virulence factors that enable *M. abscessus* to colonize the airways in humans.

## INFORMATIONS COMPLÉMENTAIRES

More information: <https://www.jove.com/video/57766/identification-virulence-markers-mycobacterium-abscessus-for>